

Quantum Electrodynamical Effects in Dusty Plasmas

M. Marklund,^{1,2} L. Stenflo,¹ P.K. Shukla,^{1,2} and G. Brodin^{1,2}

¹*Department of Physics, Umeå University, SE-901 87 Umeå, Sweden*

²*Centre for Fundamental Physics, Rutherford Appleton Laboratory,
Chilton Didcot, Oxfordshire, OX11 0QX, UK*

(Dated: February 2, 2008)

Abstract

A new nonlinear electromagnetic wave mode in a magnetized dusty plasma is predicted. Its existence depends on the interaction of an intense circularly polarized electromagnetic wave with a dusty plasma, where quantum electrodynamical photon–photon scattering is taken into account. Specifically, we consider a dusty electron–positron–ion plasma, and show that the propagation of the new mode is admitted. It could be of significance for the physics of supernova remnants and in neutron star formation.

PACS numbers: 52.27.Fp, 52.35.Mw, 52.38.-r, 52.40.Db

I. INTRODUCTION

Dusty plasmas play an important role in planetary as well as in astrophysical systems [1, 2, 3]. However, many prominent astrophysical environments in which dusty plasmas can be found are highly energetic and strongly magnetized, such as supernovae, neutron stars, and young stellar objects (YSO). There are important new wave modes in such dusty plasmas [4], and the dust grain speeds may even become relativistic [5, 6]. Supernovae, in which extreme energy scales are reached, allow for many exotic phenomena. In these violent events, neutron stars, YSO and pulsars are formed. Quantum electrodynamics (QED) is an indispensable explanatory model for much of the observed phenomena around these highly condensed objects. Scattering of photons off photons is predicted by QED, and it can be a prominent component of pulsar physics, since pulsars offer the necessary energy scales for such scattering to occur. Related to the scattering of photons is the concept of photon splitting in strong magnetic fields [7]. It has been suggested that this effect could be important in explaining the radio silence of magnetars [8, 9]. Moreover, the propagation of intense electromagnetic waves in plasmas gives rise to many interesting phenomena [10]. The existence of new plasma modes due to QED effects has been noted recently [11, 12, 13]. In the present Letter, we point out the existence of a different electromagnetic wave that may exist due to the interaction of charged dust and photons with the nonlinear quantum vacuum.

II. BASIC EQUATIONS

First, let us consider the classical propagation of a circularly polarized electromagnetic wave in a multi-component plasma embedded in an external magnetic field $B_0 \hat{\mathbf{z}}$, where B_0 is the strength of the external magnetic field and $\hat{\mathbf{z}}$ is the unit vector along the z axis. The dispersion relation for a circularly polarized wave with frequency ω and wavevector $k \hat{\mathbf{z}}$ is then given by (see e.g. Refs. [14] and [15])

$$\frac{k^2 c^2}{\omega^2} = 1 - \sum_j \frac{\omega_{pj}^2}{\omega(\omega \gamma_j \pm \omega_{cj})} \quad (1)$$

where the sum is over all particle species j , $\omega_{cj} = q_j B_0 / m_{0j}$ and $\omega_{pj} = (n_{0j} q_j^2 / \epsilon_0 m_{0j})^{1/2}$ is the gyrofrequency and plasma frequency, respectively, γ_j is the relativistic gamma factor of

species j , and the upper (lower) sign in the denominator corresponds to left (right) hand polarization. Here, n_{0j} denotes particle density in the laboratory frame, q_j charge, m_{0j} particle rest mass, and c the speed of light in vacuum. The plasma is assumed to consist of an admixture of electrons (e), positrons (p), ions (i), and charged dust particles (d). We note that the dispersion relation (1) has been derived without applying the usual small amplitude linearization. For a given k , the polarization is determined by the sign of ω . In the background state, the plasma is close to equilibrium, and we have approximate charge neutrality, $\sum_j q_j n_{0j} \approx 0$. Thus, for strongly magnetized relativistic electrons, positrons and ions, a low-frequency ($\omega \ll kc$ and $\omega\gamma_{e,p,i} \ll \omega_{ce,cp,ci}$) wave will satisfy the dispersion relation

$$\frac{k^2 c^2}{\omega^2} \approx -\frac{\omega_{pd}^2 \gamma_d}{\omega_{cd}(\mp \omega \gamma_d - \omega_{cd})}, \quad (2)$$

which is appropriate for large amplitude electromagnetic waves in supernova. Here, $\gamma_d = (1 + \nu^2)^{1/2}$, where

$$\nu^2 = \left(\frac{Z_d e E_0}{c m_{od}} \right)^2 \frac{1 + \nu^2}{[\omega(1 + \nu^2)^{1/2} \pm \omega_{cd}]^2}, \quad (3)$$

gives the relativistic gamma factor for dust with charge $q_d = Z_d e$, where e is the magnitude of the electron charge. Henceforth we will allow $\omega < 0$, thus incorporating the left hand polarization as negative frequencies, and removing the $\mp$ in Eq. (2).

Next, we investigate the effects of the nonlinear quantum vacuum on the dispersion relation (2) of a circularly polarized electromagnetic wave in a dusty magnetoplasma.

The weak field theory of photon-photon scattering can be formulated in terms of the effective Lagrangian density

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{\text{HE}}, \quad (4)$$

where $\mathcal{L}_0 = -\frac{1}{4}\epsilon_0 F_{ab}F^{ab} = \frac{1}{2}\epsilon_0(\mathbf{E}^2 - c^2\mathbf{B}^2)$ is the classical free field Lagrangian, and

$$\mathcal{L}_{\text{HE}} = \kappa \epsilon_0^2 \left[4 \left(\frac{1}{4} F_{ab} F^{ab} \right)^2 + 7 \left(\frac{1}{4} F_{ab} \hat{F}^{ab} \right)^2 \right], \quad (5)$$

is the Heisenberg-Euler correction [16, 17], where $\hat{F}_{ab} = \frac{1}{2}\epsilon_{abcd}F^{cd}$, and $\frac{1}{4}\hat{F}_{ab}F^{ab} = -c\mathbf{E} \cdot \mathbf{B}$. Here, $\kappa \equiv 2\alpha^2 \hbar^3 / 45 m_{0e}^4 c^5 \approx 1.63 \times 10^{-30} \text{ ms}^2/\text{kg}$, α is the fine-structure constant, and $\hbar$ is the Planck constant. With $F_{ab} = \partial_a A_b - \partial_b A_a$, A^b being the four-potential, we obtain (see, e.g. Ref. [18])

$$\partial_b F^{ab} = 2\epsilon_0 \kappa \partial_b \left[(F_{cd} F^{cd}) F^{ab} + \frac{7}{4} (F_{cd} \hat{F}^{cd}) \hat{F}^{ab} \right] + \mu_0 j^a, \quad (6)$$

where j^a is the four current.

For a magnetic field-aligned circularly polarized electromagnetic wave with the electric field $\mathbf{E}_0 = E_0(\hat{\mathbf{x}} + i\hat{\mathbf{y}})\exp(ikz - i\omega t)$, the invariants satisfy $F_{cd}F^{cd} = -2E_0^2(1 - k^2c^2/\omega^2) + 2c^2B_0^2$ and $F_{cd}\hat{F}^{cd} = 0$. Thus, Eq. (6) can be written as

$$\square A^a = -4\epsilon_0\kappa \left[E_0^2 \left(1 - \frac{k^2c^2}{\omega^2} \right) - c^2B_0^2 \right] \square A^a + \mu_0 j^a, \quad (7)$$

in the Lorentz gauge, and $\square = \partial_a\partial^a$. For right-hand circularly polarized electromagnetic waves propagating in a magnetized cold dusty pair-ion plasma, the matter effects can be absorbed in the wave operator on the left-hand side by the replacement $\square \rightarrow -D(\omega, k)$, where D is the plasma dispersion function, given by (see, e.g. Refs. [4, 14, 19], and Eq. (2))

$$D(\omega, k) = k^2c^2 + \frac{\omega^2\omega_{pd}^2\gamma_d}{\omega_{cd}(\omega\gamma_d - \omega_{cd})}. \quad (8)$$

We stress that the application of Eq. (5) requires field strengths smaller than the Schwinger critical field $E_S = m_{0e}^2c^3/e\hbar \sim 10^{18}$ V/m [see, e.g. Ref. [17]]. Looking for low frequency modes, the dispersion relation obtained from Eqs. (7) and (8) then reads

$$\frac{k^2c^2}{\omega^2} \approx \frac{4\alpha}{45\pi} \left[\left(\frac{E_0}{E_S} \right)^2 \frac{k^2c^2}{\omega^2} + \left(\frac{cB_0}{E_S} \right)^2 \right] \frac{k^2c^2}{\omega^2} - \frac{\omega_{pd}^2\gamma_d}{\omega_{cd}(\omega\gamma_d - \omega_{cd})}. \quad (9)$$

Equation (9) is the main result of our calculations. Due to their large mass, the dust particles can under most circumstances be considered non-relativistic, i.e. $\gamma_d \approx 1$ [5, 6]. We note the following for the dispersion relation (9): In the classical case, according to Eq. (2) we must have $\omega\gamma_d - \omega_{cd} < 0$. This requirement is obviously not necessary for the QED case, where $\omega\gamma_d - \omega_{cd} \geq 0$ is allowed. Thus, new low frequency waves are expected in the strong field regime.

III. ANALYSIS OF THE DISPERSION RELATION

In order to gain a better understanding of the dispersion relation (9) we introduce the dimensionless variables and parameters $\Omega = \omega/\omega_{cd}$, $K = kc/\omega_{pd}$, $\Omega_p = \omega_{pd}/\omega_{cd}$, $\sigma = (4\alpha/45\pi)(cB_0/E_S)^2$, and $\delta = E_0/cB_0$. The dispersion relation (9) then reads

$$\left(\frac{K}{\Omega} \right)^2 = \sigma \left[1 + \delta^2\Omega_p^2 \left(\frac{K}{\Omega} \right)^2 \right] \left(\frac{K}{\Omega} \right)^2 + \frac{\gamma_d}{1 - \Omega\gamma_d}, \quad (10)$$

where the relativistic gamma factor can be found from the relation

$$\frac{\gamma_d^2 - 1}{\gamma_d^2} = \frac{\delta^2}{(1 - \Omega\gamma_d)^2}. \quad (11)$$

We may now discern between the following cases:

(a) $\gamma_d \sim 1$: If we let $\sigma \rightarrow 0$, i.e. neglect the QED correction, we only have mode propagation for $\Omega - 1 < 0$. When $\sigma \neq 0$, this is no longer a necessary condition for mode propagation, and we can therefore have $\Omega - 1 > 0$. A special case here is the strongly magnetized dusty plasma, $cB_0 \gg E_0$. Equation (9) then yields

$$(1 - \sigma) \left(\frac{K}{\Omega} \right)^2 = \frac{1}{1 - \Omega}, \quad (12)$$

for which the QED effects act only as a small correction.

(b) $\gamma_d \gg 1$: From Eqs. (10) and (11) we obtain the dispersion relation

$$\left(\frac{K}{\Omega} \right)^2 = \sigma \left[1 + \delta^2 \Omega_p^2 \left(\frac{K}{\Omega} \right)^2 \right] \left(\frac{K}{\Omega} \right)^2 + \frac{-\delta \pm 1}{\Omega \delta}. \quad (13)$$

In the limit $\sigma \rightarrow 0$, we have a single mode given by

$$\left(\frac{K}{\Omega} \right)^2 = \frac{-\delta + 1}{\Omega \delta}. \quad (14)$$

When $\sigma \neq 0$, we have further completely distinct modes as given by the dispersion relation

$$\left(\frac{K}{\Omega} \right)^2 = \sigma \left[1 + \delta^2 \Omega_p^2 \left(\frac{K}{\Omega} \right)^2 \right] \left(\frac{K}{\Omega} \right)^2 - \frac{1 + \delta}{\Omega \delta}. \quad (15)$$

As we will demonstrate, Eq. (15) has new branches which is not present in the classical theory. Moreover, it will be further pronounced for low frequency waves, and we therefore expect the QED effects to play an important role in the propagation of such modes. In the limit of large δ , i.e. weak external magnetic field B_0 , we can simplify the dispersion relation to

$$\left(\frac{K}{\Omega} \right)^2 = \sigma \delta^2 \Omega_p^2 \left(\frac{K}{\Omega} \right)^4 - \frac{1}{\Omega}, \quad (16)$$

in order to further elucidate the character of the new QED mode. In the limit $\sigma \rightarrow 0$, we obtain the whistler-like dispersion relation $\Omega = -K^2$ (see Fig. 1).

IV. APPLICATIONS AND CONCLUSIONS

In astrophysical environments, such as supernovae, it is believed that charged dust grains can affect the plasma dynamics in crucial ways [20, 21, 22]. It has also been argued that supernovae can be a significant source of dust particles [23], although this claim is under debate [24]. At the same time the plasma may be highly magnetized, thus altering the behavior of the dusty plasma significantly [25]. In fact, the fields can reach amplitudes such that the vacuum will be excited, making it nonlinear due to QED effects. In this perspective, it is thus natural that the mode structure presented here is more complex than that of linear theory. In Fig. 1, we have plotted the dispersion curves, as given by Eq. (16), for different values of the QED parameter A . We note that for K larger than a critical value (which depends on A), there are three branches of the dispersion relation (16), in contrast to the single branch present in the classical limit. It should be stressed that the values of A used in Fig. 1 by no means require unrealistic astrophysical conditions, as the following example will show: In interstellar clouds, the dust density is expected to be $n_{0d} \sim 10^{-1} \text{ m}^{-3}$ [26], with a typical dust mass $m_{0d} \sim 10^6 m_{\text{proton}} \sim 10^{-21} \text{ kg}$ [3]. For the circularly polarized wave, we take the amplitude to satisfy $E_0/E_S \approx 2 \times 10^{-5}$, a value appropriate at a distance of 1 AU from a supernova. In this case, a value $B_0 \sim 10^{-8} \text{ G}$ (well within the limits for such environments) results in $A \sim 1$. For higher dust mass densities we note that even lower wave intensities will give values of A close to one. Such dense dusty plasmas are likely to be found closer to the core of a supernova remnant. It should also be noted that we have neglected radiative-resonant interactions on our new electromagnetic wave mode. However, in many cases such interactions can dominate [27, 28], thus altering the behavior of the mode. This is a phenomenon that has to be considered in future research. In addition, an important problem of the physics of supernova explosions is the relation between the layer of dust behind the supernova shock and the process of dust-wave condensation behind the shock wave front [29]. Our new nonlinear wave mode may thus play a role from the viewpoint of supernova shock propagation.

We have presented an analysis regarding the propagation of intense circularly polarized waves in a magnetized dusty plasma. It is found that corrections due to QED modify the mode structure, giving rise to novel low frequency waves with no classical counterpart. Our main result is the dispersion relation (9), in which the effects of charged dust and QED are

taken into account. The result should be of interest in astrophysical environments, such as supernova remnants and interstellar clouds.

Acknowledgements

This research was partially supported by the Swedish Research Council through the contracts No. 621-2001-2274 and No. 621-2004-3217, as well as by the Deutsche Forschungsgemeinschaft through the Sonderforschungsbereich 591 and the European Commission through the contract No. HPRN-CT-2001-00314.

-
- [1] M. Horányi, T.W. Hartquist, O. Havnes, D.A. Mendis, and G.E. Morfill, *Rev. Geophys.* **42**, RG4002 (2004).
 - [2] Y.K. Okamoto, H. Kataza, M. Honda, T. Yamashita, T. Onaka, J. Watanabe, T. Miyata, S. Sako, T. Fujiyoshi, and I. Sakon, *Nature* **431** 660 (2004).
 - [3] P.K. Shukla and A.A. Mamun, *Introduction to Dusty Plasma Physics* (Institute of Physics Publishing, Bristol, 2000).
 - [4] L. Stenflo and P.K. Shukla, *IEEE Trans. Plasma Sci.* **29**, 208 (2001).
 - [5] R. Bingham and V.N. Tsytovich, *Astropart. Phys.* **12** 35 (1999).
 - [6] V.N. Tsytovich and R. Bingham, *AIP Conference Proceedings* **649** 450 (2002).
 - [7] S.L. Adler, *Ann. Phys.-NY* **67** 599 (1971).
 - [8] M.G. Baring and A.K. Harding, *ApJ* **547** 929 (2001).
 - [9] C. Kouveliotou, S. Dieters, T. Strohmayer, J. van Paradijs, G.J. Fishman, C.A. Meegan, K. Hurley, J. Kommers, I. Smith, D. Frail, and T. Murakami, *Nature* **393** 235 (1998).
 - [10] J.T. Mendonça, *Theory of Photon Acceleration* (Institute of Physics Publishing, Bristol, 2001).
 - [11] M. Marklund, P.K. Shukla, L. Stenflo, G. Brodin, and M. Servin, New low-frequency nonlinear electromagnetic wave in a magnetized plasma, *Plasma Phys. Control. Fusion*, in press (2005).
 - [12] M. Marklund, P.K. Shukla, G. Brodin, and L. Stenflo, *Proceedings of the 12th International Congress on Plasma Physics, Nice (France) (25-29 October 2004)*. O3-0100; e-Proc. available on line at <http://hal.ccsd.cnrs.fr/ccsd-00003094/en> (2004b).
 - [13] L. Stenflo, G. Brodin, M. Marklund, and P.K. Shukla, A new electromagnetic wave in a pair plasma, *J. Plasma Phys.*, in press (2005).
 - [14] L. Stenflo and N.L. Tsintsadze, *Astrophys. Space Sci.* **64**, 513 (1979).
 - [15] P.K. Shukla, N.N. Rao, M.Y. Yu, and N.L. Tsintsadze, *Phys. Rep.* **138** 1 (1986).
 - [16] W. Heisenberg and H. Euler, *Z. Phys.* **98** 714 (1936).
 - [17] J. Schwinger, *Phys. Rev.* **82** 664 (1951).
 - [18] P.K. Shukla, M. Marklund, D.D. Tskhakaya, and B. Eliasson, *Phys. Plasmas* **11**, 3767 (2004).
 - [19] L. Stenflo, *Phys. Scripta* **14**, 320 (1976).
 - [20] R. Bingham and V.N. Tsytovich, *AIP Conference Proceedings* **649** 126 (2002).
 - [21] S.I. Popel and V.N. Tsytovich, *Astrophys. Space Sci.* **264** 219 (1999).

- [22] E. Sedlmayr, AIP Conference Proceedings **649** 32 (2002).
- [23] L. Dunne, S. Eales, R. Ivison, H. Morgan, and M. Edmunds, Nature **424** 285 (2003)
- [24] O. Krause, S.M. Birkmann, G.H. Rieke, D. Lemke, U. Klaas, D.C. Hines, and K.D. Gordon, Nature **432** 596 (2004).
- [25] N. Sato, AIP Conference Proceedings **649** 66 (2002).
- [26] D.A. Mendis and M. Rosenberg, Annu. Rev. Astron. Astrophys. **32** 419 (1994).
- [27] V.N. Tsytovich, Phys. Rep. **178**, 261 (1989).
- [28] V.N. Tsytovich and R. Bingham, Phys. Lett. A **259**, 393 (1999).
- [29] S.I. Popel, V.N. Tsytovich, and M.Y. Yu, Astrophys. Space Sci. **256**, 107 (1998).




FIG. 1: The normalized frequency Ω plotted as a function of the normalized wavenumber K , as given by the relation (16). Variation of the parameter $A = \sigma\delta^2\Omega_p^2$ gives the different solution curves. The curves in the upper and lower half planes represents the roots of Eq. (16). From the line $\Omega = 0$ outwards, the curves for the values $A = 0.1, 1, 2, 3$, and 4 have been plotted. We note that the modes for which $\Omega > 0$ (right-hand polarization) have no classical counterpart. When $A \rightarrow 0$ we obtain $\Omega = -K^2$, which represents a classical limit curve in the lower half plane.